

- Moore, R., Robinson, W., Lovette, I. & Robinson, T. 2008. Experimental evidence for extreme dispersal limitation in tropical forest birds. *Ecol. Lett.* 11: 960–968.
- Moores, N. 2007. Selected records from Socheong Island, South Korea. *Forktail* 23: 102–124.
- Perlman, Y. 2001. Blyth's Pipit *Anthus godlewskii*: a new species for Thailand. *Forktail* 17: 115–116.
- Pratt, H. D., Engbring, J., Bruner, P. L. & Berrett, D. G. 1980. Notes on the taxonomy, natural history, and status of the resident birds of Palau. *Condor* 82: 117–131.
- Pratt, H. D. & Etpison, M. T. 2008. *The birds and bats of Palau*. Mutual Publishing, Honolulu, HI.
- Prünte, W., Vohwinkel, R., Karaardıç, H. & Erdoğan, A. 2010. First record of Blyth's Pipit, *Anthus godlewskii* (Taczanowski, 1876), from Turkey. *Zool. Middle East* 50: 129–131.
- Pyle, P. 1984. Observations of migrant and vagrant birds from Kure and Midway atolls, 1982–1983. *'Elepaia* 44: 107–111.
- Rabøl, J. 1976. The orientation of Pallas's Leaf Warbler *Phylloscopus proregulus* in Europe. *Dansk. Orn. Foren. Tidsskr.* 70: 6–16.
- Richardson, C. 2008. *Cyprus Bird Report 2007*. BirdLife Cyprus, Cyprus.
- Slack, R. 2009. *Rare birds, where and when: an analysis of status and distribution in Britain and Ireland*, vol. 1. Rare Bird Books, York.
- VanderWerf, E. A. 2007. 2005 bird surveys in the Republic of Palau, final report. Unpubl. rep. to US Fish & Wildlife Service and Palau Conservation Society. Pacific Rim Conservation, Honolulu, HI.
- Vanderwerf, E. A., Wiles, G. J., Marshall, A. P. & Knecht, M. 2006. Observations of migrants and other birds in Palau, April–May 2005, including the first Micronesian record of a Richard's Pipit. *Micronesica* 39: 11–29.
- Wernham, C. V., Toms, M. P., Marchant, J. H., Clarke, J. A., Siriwardena, G. M. & Baillie, S. R. 2002. *The migration atlas: movements of the birds of Britain and Ireland*. T. & A. D. Poyser, London.
- Wiles, G. J. 2005. A checklist of the birds and mammals of Micronesia. *Micronesica* 38: 141–189.
- Williamson, K. 1977. Blyth's Pipit in the Western Palearctic. *Bull. Brit. Orn. Cl.* 97: 60–61.
- Addresses:* Alexander C. Lees, Dept. of Zoology, Museu Paraense Emilio Goeldi, C.P. 399, CEP 66040-170, Belém, Pará, Brazil, e-mail: alexanderlees@btopenworld.com. EricVanderWerf, Pacific Rim Conservation, 3038 Oahu Avenue, Honolulu, Hawaii 96822, USA, e-mail: ewerf@hawaii.rr.com

© British Ornithologists' Club 2011

Expansion of the known range of Tawny Piculet *Picumnus fulvescens* including the south bank of the São Francisco River in north-east Brazil

by Juan Ruiz-Esparza, Patrício Adriano da Rocha, Adauto de Souza Ribeiro,
Stephen F. Ferrari & Helder F. P. Araujo

Received 28 April 2011

Tawny Piculet *Picumnus fulvescens* is a small-bodied Picidae, with predominantly tawny-coloured plumage and ferruginous, white-speckled underparts. Males are distinguished from females by the red forehead (Sick 1997). The species is currently listed as Near Threatened, due primarily to extensive and ongoing habitat loss within its range (BirdLife International 2008). The species occurs predominantly in Caatinga scrub, but also in drier forests of the Atlantic Forest domain. Like other piculets, *P. fulvescens* appears relatively tolerant of habitat disturbance, and is often found in second growth, where it feeds on insects, particularly the larvae and pupae of ants (Sick 1997).

P. fulvescens is endemic to north-east Brazil, in the northern Caatinga east to the coastal Atlantic Forest. Prior to the present study, the species' known range (*cf.* Ridgely *et al.* 2007) encompassed a relatively limited area parallel to the left (north) bank of the São Francisco River in the states of Alagoas, Ceará, Pernambuco and south-east Piauí (Fig. 1).




Figure 1. Distribution of Tawny Piculet *Picumnus fulvescens* in north-east Brazil. The grey shaded area is the range defined by Ridgely *et al.* (2007), with localities marked by white circles. Stars represent new localities recorded by the present study and triangles sites mentioned in the literature or based on the Federal University of Pernambuco (UFPE), Recife, ornithology collection. Sites numbered as in Table I. Brazilian states: AL = Alagoas; BA = Bahia; CE = Ceará; MG = Minas Gerais; PB = Paraíba; PE = Pernambuco; PI = Piauí; RN = Rio Grande do Norte; SE = Sergipe.

During field work at the Serra da Guia in western Sergipe, an adult male *P. fulvescens* (Fig. 2) was captured in a mist-net at 17.00 h on 19 December 2008. The trapping site (09°58'55"S, 37°52'06"W) is located within an enclave of cloud forest in an area of arboreal caatinga, 750 m above sea level. In October 2010, the species was observed during field work at a neighbouring site in western Sergipe (09°45'22"S, 37°51'12"W), 430 m above sea level.

TABLE 1

Geographic localities identified for Tawny Piculet *Picumnus flavescens* during the present study. Brazilian states listed by locality: AL = Alagoas; CE = Ceará; PB = Paraíba; PE = Pernambuco; RN = Rio Grande do Norte; SE = Sergipe. Habitats: CA = Caatinga; AF = Atlantic Forest.

Site	Locality	Coordinates	Habitat	Source
1	Piaçabuçu (AL)	10°16'S, 36°22'W	AF	Cabral <i>et al.</i> (2006)
2	Murici (AL)	09°19'S, 35°56'W	AF	Roda (2003)
3	Quebrangulo (AL)	09°19'S, 36°28'W	AF	Roda (2003)
4	São José da Laje (AL)	09°01'S, 36°03'W	AF	Roda (2003)
5	Palmares (PE)	08°41'S, 35°35'W	AF	Roda (2003)
6	Buíque (PE)	08°27'S, 37°15'W	CA	Farias (2009)
7	Brejão (PE)	09°01'S, 36°34'W	CA	Roda & Carlos (2004)
8	Floresta (PE)	08°36'S, 35°34'W	AF	UFPE specimen
9	Sertânia (PE)	08°58'S, 38°13'W	CA	UFPE specimen
10	Caruaru (PB)	08°17'S, 38°13'W	CA	UFPE specimen
11	Sertânia (PE)	08°04'S, 37°15'W	CA	UFPE specimen
12	São Vicente Ferrer (PE)	07°35'S, 35°26'W	AF	Roda & Carlos (2004)
13	Betânia (PE)	08°18'S, 38°12'W	CA	Farias <i>et al.</i> (2006)
14	Icó Mandante (PE)	08°52'S, 38°17'W	CA	Farias (2007)
15	Brígida (PE)	08°28'S, 39°32'W	CA	Farias (2007)
16	Apolônio Sales (PE)	08°54'S, 38°14'W	CA	Farias (2007)
17	Coremas (PB)	07°03'S, 37°53'W	CA	Pinto & Camargo (1961)
18	Santa Teresinha (PB)	07°01'S, 37°24'W	CA	Telino-Júnior <i>et al.</i> (2005)
19	Curimataú (PB)	06°40'S, 35°45'W	CA	Farias <i>et al.</i> (2006)
20	Areia (PB)	06°58'S, 35°44'W	CA	H. F. P. Araujo pers. obs.
21	Mataraca (PB)	06°33'S, 34°58'W	AF	Araujo <i>et al.</i> (2010)
22	Mamanguape (PB)	06°43'S, 35°10'W	AF	Almeida & Teixeira (2010)
23	Sumé (PB)	07°40'S, 36°57'W	CA	Araujo (2009)
24	Maturéia (PB)	07°15'S, 37°12'W	CA	H. F. P. Araujo pers. obs.
25	São José da Lagoa Tapada (PB)	06°57'S, 38°10'W	CA	H. F. P. Araujo pers. obs.
26	Serra Negra (RN)	06°34'S, 37°30'W	CA	Nascimento (2000)
27	Chapada do Araripe (CE)	07°14'S, 39°56'W	CA	Nascimento <i>et al.</i> (2000)
28	Pedra Branca (CE)	05°40'S, 39°38'W	CA	Olmos <i>et al.</i> (2005)
29	Mombaça (CE)	05°40'S, 39°41'W	CA	Olmos <i>et al.</i> (2005)
30	Milagres (CE)	07°22'S, 38°58'W	CA	Olmos <i>et al.</i> (2005)
31	Jati (CE)	07°42'S, 39°01'W	CA	Olmos <i>et al.</i> (2005)
32	Serra da Guia (SE)	09°59'S, 37°52'W	CA	Present study
33	Capim Grosso (SE)	09°45'S, 37°51'W	CA	Present study

As these records represent a considerable expansion of the species' known range, we reviewed the literature and consulted specimens in the ornithological collection of the Federal University of Pernambuco (UFPE) in Recife (Table 1). As a result, we identified an additional 28 localities for *P. fulvescens*, extending the limits of its known range in almost all directions, except west and north-west (Fig. 1). Three of these 'new' localities are on the left (north) bank of the São Francisco River in Alagoas and Pernambuco.

In addition to confirming that *P. fulvescens* occurs over a much wider area than previously thought (Ridgely *et al.* 2007), our records from Sergipe confirm its presence south of the São Francisco River, a major physical barrier to small-bodied forest-dwelling birds (Welty & Baptista 1988). It is therefore possible that the species is more widespread in Sergipe and even Bahia, although the current lack of records may at least partially reflect its

rarity therein (Wege & Long 1995, BirdLife International 2008).

The records reported here support the recent reclassification of *P. fulvescens* as Near Threatened rather than Vulnerable (BirdLife International 2008). In fact, this review suggested that the species may even warrant downlisting to Least Concern, and our study would support this, given the considerable expansion of this piculet's range. However, while the species is now known to occur over a relatively wide area, including a number of protected areas, Tawny Piculet does appear to be relatively rare throughout its distribution.



Figure 2. Adult male Tawny Piculet *Picumnus fulvescens*, Serra da Guia, Sergipe, Brazil, 19 December 2008 (Juan Ruiz-Esparza)

Acknowledgements

This study was supported by the Sergipe state environment ministry (SEMARTH), FAPITEC-Sergipe, CAPES and CNPq (project nos. 302747/2008-7 and 476064/2008-2). We are especially grateful to Zefa da Guia, and we thank also Raone Beltrão, Eduardo Marques, Juliana Cordeiro, Douglas Matos, Thiago Bicudo, Sidney Gouvêia and Validineide Santana for their practical help. We are grateful to Weber Silva and Sidnei Dantas for their helpful comments on the submitted version of this manuscript.

References:

- Almeida, A. C & Teixeira, D. 2010. Aves da Reserva Biológica Guaribas, Mamanguape, Paraíba, Brasil. *Rev. Nordestina Biol.* 19: 3–14.
- Araujo, H. F. P. 2009. Amostragem, estimativa de riqueza de espécies e variação temporal na diversidade, dieta e reprodução de aves em área de caatinga, Brasil. Ph.D. thesis. Universidade Federal de Paraíba, João Pessoa.
- Araujo, H., Mariano, E., Toledo, G., Vieira Filho, A. & Hernández, M. 2010. Avifauna de floresta de restinga em um complexo de mineração no litoral norte da Paraíba, Brasil. *Rev. Nordestina Zool.* 4: 46–56.
- BirdLife International. 2008. *Picumnus fulvescens*. In IUCN. 2010. IUCN Red List of threatened species. Version 2010.4. www.iucnredlist.org.
- Cabral, S. A. S., Azevedo Júnior, S. M. & Larrazábal, M. E. 2006. Levantamento de aves da Área de Proteção Ambiental de Piaçabuçu, no litoral de Alagoas, Brasil. *Ornithologia* 1: 161–167.
- Farias, G. B. 2007. Avifauna em quatro áreas de caatinga *strictu sensu* no centro-oeste de Pernambuco, Brasil. *Rev. Bras. Orn.* 15: 53–60.
- Farias, G. B. 2009. Aves do Parque Nacional Pernambuco, Brasil. *Atualidades Orn.* 147: 29 [www.ao.com.br].
- Farias G. B., Girão e Silva, W. A. & Albano, C. G. 2006. Diversidade de aves em áreas prioritárias para conservação de aves da Caatinga. Pp. 204–226 in Araujo, F. S., Rodal, M. J. N. & Barbosa, M. R. V. (eds.) *Análise das variações da biodiversidade do bioma caatinga. Suporte a estratégias regionais de conservação*. Ministério do Meio Ambiente, Brasília.
- Nascimento, J. L. X. 2000. Estudo comparativo em duas Estações Ecológicas da Caatinga: Aiuaba e Seridó. *Melopsittacus* 3: 12–35.
- Nascimento, J. L. X., Nascimento, L. S. & Azevedo-Júnior, S. M. 2000. Aves da Chapada do Araripe (Brasil): biologia e conservação. *Ararajuba* 8: 115–125.
- Olmos, F., Girão e Silva, W. A. & Albano, C. G. 2005. Aves de oito áreas de Caatinga no sul do Ceará e oeste de Pernambuco, Nordeste do Brasil: composição, riqueza e similaridade. *Pap. Avuls. Zool., São Paulo* 45: 179–199.
- Pinto, O. M. O. & Camargo, E. A. 1961. Resultados ornitológicos de quatro recentes expedições do Departamento de Ornitologia ao nordeste do Brasil, com a seis nova subespécies. *Arq. Zool., São Paulo* 11: 193–284.
- Ridgely, R. S., Allnutt, T. F., Brooks, T., McNicol, D. K., Mehlman, D. W., Young, B. E. & Zook, J. R. 2007. Digital distribution maps of the birds of the Western Hemisphere, version 3.0. NatureServe, Arlington, VA.
- Roda, S. 2003. Aves do Centro de Endemismo Pernambuco: composição, biogeografia e conservação. Ph.D. thesis. Universidade Federal do Pará, Belém.

- Roda, S. A. & Carlos J., C. J. 2004. Composição e sensibilidade da avifauna dos brejos de altitude do Estado de Pernambuco. Pp. 211–228 in Porto, K., Cabral, J. & Tabarelli, M. (eds.) *Brejos de altitude: história natural, ecologia e conservação*. Ministério do Meio Ambiente, Brasília.
- Sick, H. 1997. *Ornitologia brasileira*. Ed. Nova Fronteira, Rio de Janeiro.
- Telino-Júnior, W. R., Lyra-Neves, R. M. & Nascimento, J. L. X. 2005. Biologia e composição da avifauna em uma Reserva Particular de Patrimônio Natural da caatinga paraibana. *Ornithologia* 1: 49–57.
- Wege, D. C. & Long, A. J. 1995. *Key Areas for threatened birds in the Neotropics*. BirdLife International, Cambridge, UK.
- Welty, J. C. & Baptista, L. 1988. *The life of birds*. Fourth edn. Saunders College Publishing, New York.

Addresses: Juan Ruiz-Esparza, Graduate Program in Ecology and Conservation, and PRODEMA, Universidade Federal de Sergipe, Av. Marechal Rondon s/n, 49.100-000 São Cristóvão, Sergipe, Brazil, e-mail: juancolorado21@hotmail.com. Patrício Adriano da Rocha, Graduate Program in Zoology, Universidade Federal da Paraíba, João Pessoa, Paraíba, Brazil. Adauto de Souza Ribeiro & Stephen F. Ferrari, Dept. of Biology, Universidade Federal de Sergipe, São Cristóvão, Sergipe, Brazil. Helder F. P. Araujo, Dept. of Biological Sciences, Universidade Federal da Paraíba, João Pessoa, Paraíba, Brazil.

© British Ornithologists' Club 2011

The availability and validity of the name *Forpus flavicollis* Bertagnolio & Racheli, 2010, for a parrotlet from Colombia

David G. Notton

Received 19 May 2011

The recent description of the Yellow-necked Parrotlet *Forpus flavicollis* Bertagnolio & Racheli, 2010, from a photograph taken in Colombia has generated extensive discussion among ornithologists about methods of description through illustrations and lack of deposited type specimens. This note aims to clarify these issues on the availability and validity for this scientific name.

In zoological nomenclature there is an important distinction in the meaning of the technical terms 'availability' and 'validity'. *Available* names are those which must be taken into account as a part of zoological nomenclature. Names that are *not available* effectively do not exist for the purposes of zoological nomenclature; they cannot enter into synonymy or homonymy, and cannot be used as the names of taxa. In contrast, a *valid name* is the name that is currently thought to be the correct name for a taxon, i.e. the oldest *available name* of a name-bearing type specimen that falls with an author's concept of the taxon. Consequently, a name may be available but not valid, e.g. a junior synonym. A valid name must be one selected from the available names in zoological nomenclature. A species should have only one valid name but may have many invalid names, e.g. as junior synonyms. This should not be confused with the technical use of the word valid when applied to publications, when it means the work complies with Art. 8 of the Code (ICZN 1999).

After examining the evidence, it appears that the name *F. flavicollis* is *available* in the technical sense for zoological nomenclature. It meets the requirements of the ICZN Code for establishing a new name. However the *validity* of the name remains a decision for taxonomists. They must decide whether the recognition of this species is justified or not, based on their assessment of taxonomic evidence and their taxonomic philosophy. If it is not felt that this name represents a *valid* species, then the name is a synonym or the taxon may